



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

112. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

In *Hall and Knight's Higher Algebra*, I find the following:

If $a+b+c=0$, then

$$\frac{a^5+b^5+c^5}{5} = \frac{a^3+b^3+c^3}{3} \cdot \frac{a^2+b^2+c^2}{2}; \quad \frac{a^7+b^7+c^7}{7} = \frac{a^5+b^5+c^5}{5} \cdot \frac{a^2+b^2+c^2}{2};$$

also if $a+b+c+d=0$, then

$$\frac{a^5+b^5+c^5+d^5}{5} = \frac{a^3+b^3+c^3+d^3}{3} \cdot \frac{a^2+b^2+c^2+d^2}{2}.$$

QUERY. Is there a general law governing such expressions? Investigate.

*** Solutions of these problems should be sent to J. M. Colaw not later than January 10.

GEOMETRY.

131. Proposed by J. W. YOUNG, Graduate Student, Ohio State University, Columbus, Ohio.

Prove that $\lambda + \mu\omega + \nu\omega^2$, where λ, μ, ν are integers whose sum is ± 1 , represents the points of a quilt formed by regular hexagons. ω = primitive cube root of unity. [From *Harkness and Morley's Introduction to Theory of Functions*.]

132. Proposed by ELMER SCHUYLER, Reading, Pa.

To draw a circle to cut two given circles orthogonally.

133. Proposed by P. C. CULLEN, Principal of Public Schools, Indianola, Neb.

If the two bisectors, trisectors, quadrisectors, etc., of the base angles of a triangle are mutually equal, show that the triangle is isosceles.

134. Proposed by J. C. GREGG, A. M., Superintendent of Schools, Brazil, Ind.

If $ABCD$ is a quadrilateral circumscribing a circle, show that the line joining the middle points of the diagonals AB, CD passes through the center of the circle.

*** Solutions of these problems should be sent to B. F. Finkel not later than January 10.

CALCULUS.

102. Proposed by G. B. M. ZERR, A. M., Ph.D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

A right cone has its vertex at the focus of a paraboloid of revolution, the axis of the cone perpendicular to the axis of the paraboloid. Find the volume common to both.

103. Proposed by HON. JOSIAH H. DRUMMOND, LL. D., Portland, Me.

A park, in the shape of an ellipse whose diameters are 100 and 50 rods, respectively, is surrounded by a wall: one end of a rope, whose length is the circumference of the ellipse, is fastened (outside of the wall) at one end of the longer diameter and the other end at the other end of the same diameter. Over how much surface will a horse graze, which is fastened to a ring moving freely on the rope?

*** Solutions of these problems should be sent to J. M. Colaw not later than January 10.